

SHORT REPORT

Quantifying the resilience of plant communities under different grazing intensities in a degraded shrubland: A case study in Mandalgobi, Mongolia

Takehiro Sasaki¹, Tomoo Okayasu¹, Yasuhito Shirato², Jamsran Undarmaa³ and Kazuhiko Takeuchi¹

¹ Department of Ecosystem studies, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

² National Institute for Agro-Environmental Sciences, 3-1-3 Kannondai, Tsukuba, Ibaragi 305-8064, Japan

³ Center for Ecosystem Study, Mongolian State University of Agriculture, 210153 Ulaanbaatar, Mongolia

Keywords: Arid and semiarid rangeland; plant functional types; positive feedback.

Received: 28 November 2006; accepted: 13 March 2007

Correspondence: Takehiro Sasaki, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan. Email: aa67123@mail.ecc.u-tokyo.ac.jp

doi: 10.1111/j.1744-697X.2007.00088.x

Introduction

Resilience is the changing rate at which a system returns to its former state after it has been perturbed from that state (Gunderson 2000). Arid and semiarid ecosystems are vulnerable ecosystems; losses of resilience make the system more fragile (van de Koppel *et al.* 1997; Scheffer *et al.* 2001). Most field studies on the relationship between vegetation and grazing in arid rangelands have shown that the quantity of vegetation or the species composition changes according to grazing intensities (e.g. Fernandez-Gimenez and Allen-Diaz 2001; Sasaki *et al.* 2005). However, few studies have measured changes in vegetation and grazing intensities to quantify the resilience.

Plant functional types (PFT) have been widely used by ecologists to quantify the relationship between vegetation and various disturbances (e.g. Lavorel *et al.* 1997; Sternberg *et al.* 2000) because PFT are nonphylogenetic groupings of species that show close similarities in their response to environmental and biotic controls such as grazing (Smith *et al.* 1997). Previous rangeland studies have suggested that the responses of a community to grazing are strongly associated with plant life-form and palatability to livestock (Sternberg *et al.* 2000; Fernandez-Gimenez and Allen-Diaz 2001).

In the present study, we quantified the resilience of plant communities under different grazing intensities in a degraded shrubland in southern Mongolia. We hypothesized that the community resilience in a particular site differs according to the grazing intensity.

Materials and methods

Study area and experimental design

The study was conducted in Mandalgobi, Dundgobi Province, Mongolia (45°47'N, 106°11'E). From 1993–2001, the mean annual air temperature was approximately 2°C and the total annual precipitation ranged 150–225 mm (data provided by the Institute of Meteorology and Hydrology, Ministry of Nature and Environment, Mongolia).

In the study area, there is a livestock camp (hereafter, the camp) where nomadic people start to graze their livestock and this is located approximately 10–15 km away from the town of Mandalgobi. Most livestock grazed at this area are sheep (~500 head). The soil texture (0.0–5.0 cm) is sand (Sasaki *et al.* in press), and the landform consists of gently sloping hills (generally less than 1.5°). Nomenclature follows Grubov (1982).

We assumed that the grazing intensity would decrease radially outwards from the camp. In summer 2004, we performed a preliminary survey and determined the changes in vegetation at increasing distances from the camp. Distance was measured by using a global positioning system (GPS). The area surrounding the camp was then stratified into three impact zones: <100 m, 100–400 m and >400 m from the camp. Based on this stratification, three exclosures were established for restoration at the distances of 75 m (site 1), 150 m (site 2) and 550 m (site 3) in summer 2004. Sizes of each exclosure were 15 m × 25 m, 20 m × 40 m, and 50 m × 100 m, respectively. We acknowledged that this design was in fact a pseudo-replicated design. However, our decision to establish

only one enclosure as large as possible in each impact zone rather than ensuring the statistical replicates by setting small cages allowed us to account for the potential spatial heterogeneity of vegetation pattern and process based on our knowledge of the study area.

Sampling design

We performed a vegetation survey in July 2005. We set 10 quadrats in the open and enclosure area of each site (2 m × 2 m at site 3, 1 m × 1 m at sites 1 and 2, based on the patch size of *Caragana microphylla*). These quadrats were located systematically, taking into consideration the spatial heterogeneity of vegetation pattern and process. We therefore accounted these samples to be statistical replicates in the following analysis. In each quadrat, we recorded the cover (%) and the height (cm) of all plant species.

Soil was sampled from each open site (six replicates, at a depth of 0.0–10.0 cm) in October 2004. In the laboratory, we analyzed the soil organic carbon (SOC) using a dry combustion method (Matejovic 1993).

Data analysis

We calculated the volume of each plant species in each quadrat by multiplying the cover and height of each plant species (i.e. vegetation volume; *sensu* Ohtuka *et al.* 1993; Kawada *et al.* 2005).

We used a plant functional approach (Sternberg *et al.* 2000) to quantify the differences in the resilience of plant communities on the assumption that perennial herbs (or palatable

species) were more abundant in a mixed community with shrubs in sandy soils (Gunin *et al.* 1999) before the community was perturbed. The 24 plant species found in the survey were classified by PFT (which we defined by life history and relative palatability). Relative palatability of plants was classified according to the interviews with nomadic people, the information from Mongolian botanists, and *Forage Plants of Mongolia* (Jigjidsuren and Johnson 2003). We analyzed the variation in the individual quadrats using two-way ANOVA to examine the factors grazing intensity (three levels), enclosure (two levels) and their interactions. In the ANOVA model, we used the vegetation volume of each PFT as a parameter. If the interaction was significant, then a Tukey post-hoc comparison was performed. We also explored the differences in soil organic carbon according to grazing intensities using Bonferroni multiple comparison tests. All statistical analyses were performed using R (version 2.1.1, R Development Core Team 2005).

Results

Effects of grazing intensity and enclosure on the abundance of plant functional types

Results of the two-way ANOVA for PFT revealed that shrubs (e.g. *C. microphylla*; Figure 1a) and moderately palatable species (e.g. *C. microphylla* and *Convolvulus ammannii*; Figure 1e) were affected significantly only by grazing intensity (Table 1). In both perennial herbs (e.g. *Stipa krylovii*, *Agropyron cristatum* and *Cymbaria dahurica*; Figure 1b) and palatable




Figure 1 Effects of grazing intensity and enclosure treatments on the vegetation volume of each plant functional type among the sites. (a) Shrubs, (b) perennial herbs, (c) annual herbs, (d) palatable species, (e) moderately palatable species and (f) low-palatable species. Bars represent open sites (■) and enclosure sites (□). Vegetation volume was calculated by multiplying the cover (%) and height (cm). If the interaction between grazing intensity and enclosure was significant (see also Table 1), Tukey post-hoc comparison was performed. Dissimilar letters above bars denote statistically significant differences ($P < 0.05$).

Table 1 Results from the two-way ANOVA of plant functional types (PFT)

Treatment	d.f.	Life-form						Palatability					
		Shrubs		Perennial herbs		Annual herbs		Palatable species		Moderately-palatable species		Low-palatables species	
		F	P	F	P	F	P	F	P	F	P	F	P
Grazing intensity	2	3.818	0.030	33.912	<0.001	0.652	0.526	36.881	<0.001	3.820	0.030	0.670	0.517
Exclosure	1	0.016	0.900	9.162	0.004	3.637	0.063	12.717	<0.001	0.016	0.899	0.868	0.357
Grazing intensity × Exclosure	2	0.093	0.911	6.247	0.004	0.675	0.514	5.746	0.006	0.092	0.913	0.270	0.765

Bold figures indicate significant differences ($P < 0.05$). In the model, we used a vegetation volume of each PFT as a parameter. Treatments included manipulations of grazing intensity (sites 1, 2 and 3), exclosure (open sites versus exclosure sites), and their interaction.

species (e.g. *S. krylovii* and *A. cristatum*; Figure 1d) there was a significant interaction between grazing intensity and exclosure (Table 1); the effect of exclosure was evident in the less grazed site (site 3). Although the vegetation volume of annual herbs (e.g. *Heteropappus biennis*) increased in the exclosure of sites 2 and 3 (Figure 1c), and low-palatable species (e.g. *Panzeria lanata*) decreased in all exclosures (Figure 1f), the results of the two-way ANOVA showed that grazing intensity and exclosure did not have significant effects on these PFT (Table 1).

Changes in soil organic carbon according to grazing intensities

Changes in the soil organic carbon among the sites are shown in Figure 2. The soil organic carbon contents in sites 1 and 2 were significantly lower than that in site 3 ($P < 0.05$).

Discussion

A key finding of the restorative experiment was that PFT such as perennials and palatable species did not recover in




Figure 2 Soil organic carbon (SOC; %) at sites 1, 2 and 3. SOC (+ SD) at depths of 0.0–10.0 cm ($n = 6$) are shown. Dissimilar letters above bars denote statistically significant differences ($P < 0.05$, Bonferroni multiple comparison tests). Note that the wind erosion rate is greater at sites 1 and 2.

abundance at sites 1 and 2, whereas these PFT did recover at site 3 (Figure 1b,d; Table 1). Resilience of a particular site may differ from that of another site even at a local scale. SOC decreased at sites 1 and 2; this may be evidence that wind erosion is greater at sites 1 and 2 (Figure 2), as wind erosion can cause losses of SOC (Yan *et al.* 2005). Around sites 1 and 2, vegetation volumes of PFT were generally less than those at site 3 (Figure 1), indicating potential positive feedbacks between vegetation cover and wind erosion. Previous studies demonstrated that vegetation plays an important role in protecting topsoil against wind erosion, as the presence of vegetation decreases wind force exerted on the soil surface (Buckley 1987). In contrast, wind erosion fractures bare vegetation cover mainly due to wind stream damage to plants and soil nutrient losses (Zhao *et al.* 2005). The difficulty of plant recovery at sites 1 and 2 was consistent with previous studies that reported positive feedbacks between biotic factors and components of the physical environment (van de Koppel *et al.* 1997; Suding *et al.* 2004).

At site 3, however, PFT such as perennial herbs and palatable species did recover in abundance (Figure 1b,d; Table 1). *C. microphylla* dominated both with greater size and higher abundance at site 3 than at sites 1 and 2 (T. Sasaki, unpublished data). In arid and semiarid areas, large plants may have a positive effect on understory species (Raffaele and Veblen 1998), and shrubs such as *C. microphylla* generally reduce the intensity of wind erosion (Su *et al.* 2004). Nurse effects (Raffaele and Veblen 1998) of *C. microphylla* as a benefactor species may possibly limit the positive feedbacks and support the resilience at site 3. Moreover, *C. microphylla* gathers sand around its mound, with the depth of sand deposition around each plant apparently depending on canopy size and volume (T. Sasaki, personal observation). Sand burial often has a positive effect on stabilization during seedling establishment, stimulation of growth and flowering (Maun 1997). Katoh *et al.* (1998) also reported that *C. microphylla* may play a role as a nurse plant; however, because sheep like to forage *Caragana* leaves, this species will

not act as a nurse plant under heavy grazing intensities. In the study area, most livestock are sheep and the relative palatability of *C. microphylla* is moderately high. Nurse effects of *C. microphylla* may therefore be much less at heavily grazed sites.

Although the experimental period was short, differences in the community resilience were relatively clear. However, long-term monitoring is needed to more definitively examine the community resilience that may underlie the vegetation dynamics. Vegetation responded to intense grazing with a decrease of shrubs, perennial herbs, palatable species and moderately palatable species. These results essentially agree with those of the previous studies (e.g. Naveh and Whittaker 1979; Newsome and Noble 1986).

The present study revealed differences in resilience at a local scale. After grazing removal, grazing-sensitive PFT did not recover in abundance in a formerly heavily grazed site, whereas these PFT did recover in abundance at a lightly grazed site. Our findings may provide additional empirical evidence for ecological theories of resilience.

Acknowledgments

We thank the project members, especially A. Mori and Y. Yamaguchi for their kindness in helping with the fieldwork. The staff of the Laboratory of Landscape Ecology and Planning, the University of Tokyo (especially S. Okubo and S. Yamada) provided valuable suggestions. Two anonymous referees contributed significantly to the clarity of the manuscript. This work was carried out under a JSPS research fellowship to T. Sasaki, with additional support from the Global Environmental Research Fund of the Ministry of the Environment, Government of Japan.

References

- Buckley R (1987) The effect of sparse vegetation on the transport of dune sand by wind. *Nature* 325: 426–428.
- Fernandez-Gimenez M, Allen-Diaz B (2001) Vegetation change along gradients from water sources in three grazed Mongolian ecosystems. *Plant Ecol* 157: 101–118.
- Grubov VI (1982) *Key to the Vascular Plants of Mongolia*. Science Publishers, Enfield, New Hampshire, USA.
- Gunderson LH (2000) Ecological resilience: in theory and application. *Annu Rev Ecol Syst* 31: 425–439.
- Gunin PD, Vostokova EA, Dorefeyuk NI, Tarasov PE, Black CC (1999) *Vegetation Dynamics of Mongolia*. Kluwer Academic Publishers, Boston, USA.
- Jigjidsuren S, Johnson DA (2003) *Forage Plants of Mongolia*. Admon Publishing Co, Ulaanbaatar, Mongolia.
- Katoh K, Takeuchi K, Jiang D, Nan Y, Kou Z (1998) Vegetation restoration by seasonal enclosure in the Kerqin Sandy Land, Inner Mongolia. *Plant Ecol* 139: 133–144.
- Kawada K, Vovk AG, Filatova OV, Araki M, Nakamura T, Hayashi I (2005) Floristic composition and plant biomass production of steppe communities in the vicinity of Kharkiv, Ukraine. *Grassl Sci* 51: 205–213.
- van de Koppel J, Rietkerk M, Weissing FJ (1997) Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends Ecol Evol* 12: 352–356.
- Lavorel S, McIntyre S, Landsberg J, Forbes TDA (1997) Plant functional classifications: from general groups to specific groups based on response to disturbance. *Trends Ecol Evol* 12: 474–478.
- Matejovic I (1993) Determination of carbon, hydrogen, and nitrogen in soils by automated elemental analysis (dry combustion method). *Commun Soil Sci Plan* 24: 2213–2222.
- Maun MA (1997) Adaptations of plants to burial in coastal sand dunes. *Can J Bot* 76: 713–738.
- Naveh Z, Whittaker RH (1979) Structural and floristic diversity of shrublands and woodlands in northern Israel and other Mediterranean areas. *Vegetatio* 41: 171–190.
- Newsome AE, Noble IR (1986) Ecological and physiological characters of invading species. In: *Ecology of Biological Invasions: an Australian Perspective* (eds Groves RH, Burdon JJ). Australian Academy of Sciences, Canberra, 1–20.
- Ohtuka T, Sakura T, Ohsawa T (1993) Early herbaceous succession along a topographical gradient on forest clear-felling sites in mountainous terrain, central Japan. *Ecol Res* 8: 329–340.
- R Development Core Team (2005) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Raffaello E, Veblen TT (1998) Facilitation by nurse shrubs of resprouting behavior in a post-fire shrubland in northern Patagonia, Argentina. *J Veg Sci* 9: 693–698.
- Sasaki T, Okayasu T, Takeuchi K, Jamsran U, Jadambaa S (2005) Patterns of floristic composition under different grazing intensities in Bulgan, South Gobi, Mongolia. *Grassl Sci* 51: 235–242.
- Sasaki T, Okayasu T, Shirato Y, Jamsran U, Okubo S, Takeuchi K (in press) Can edaphic factors demonstrate landscape-scale differences in vegetation responses to grazing? *Plant Ecol*.
- Scheffer M, Carpenter S, Foley JA, Folke C, Walker B (2001) Catastrophic shifts in ecosystems. *Nature* 413: 591–596.
- Smith TM, Shugart HH, Woodward FI (1997) *Plant Functional Types: Their Relevance to Ecosystem Properties and Global Change*. Cambridge University Press, Cambridge.
- Sternberg M, Gutman M, Perevolotsky A, Ungar ED, Kigel J (2000) Vegetation response to grazing management in a Mediterranean herbaceous community: a functional group approach. *J Appl Ecol* 37: 224–237.
- Su YZ, Zhao HL, Li YL, Cui JY (2004) Influencing mechanisms of several shrubs on soil chemical properties in semiarid Horqin Sandy Land, China. *Arid Land Res Manag* 18: 251–263.
- Suding KN, Gross KL, Houseman GR (2004) Alternative states and positive feedbacks in restoration ecology. *Trends Ecol Evol* 19: 46–53.
- Yan H, Wang SQ, Wang CY, Zhang GP, Patel N (2005) Losses of soil organic carbon under wind erosion in China. *Global Change Biol* 11: 828–840.
- Zhao HL, Zhao XY, Zhou RL, Zhang TH, Drake S (2005) Desertification processes due to heavy grazing in sandy rangeland, Inner Mongolia. *J Arid Environ* 62: 309–319.