

3.14

For Treatment of constipation

BEAULUCK

(Cautions)

Followings should be complied to avoid aggravation of the symptoms or incidence of adverse reactions or accidents.

1. During the treatment with BEAULUCK, do not take other laxatives.
2. Do not take overdose of BEAULUCK.

Following persons should consult doctors.

- (1) Those who are being treated by the doctor.
- (2) Women who are pregnant or are considered to be pregnant.
- (3) Persons who have symptoms of severe abdominal pain, nausea and vomiting.

When encountered with the following symptoms, discontinue immediately medication and consult doctors showing the package insert.

- (1) Following symptoms which appear right after taking medicine.

<u>Location</u>	<u>Symptoms</u>
Digestive system	Severe abdominal pain, nausea, vomiting

- (2) When symptoms are not improved in spite of continued medication for 1 week.

When diarrhea continue or be aggravated, discontinue medication and consult doctors or dentists.

(Ingredients)

Each 3 tablets (daily dose) contains Bisacodyl[4,4'-(pyridin-2-ylmethylene)bis(phenyl acetate)].....15mg

Lactose Hydrate, Corn Starch, Hydroxypropyl cellulose, Crospovidone, Methacrylic Acid Copolymer LD, Triethyl Citrate, Talc, Refined Sugar, Powdered Acacia, Gelatin, Precipitated Calcium Carbonate, Titanium Oxide, Macrogol, Carmellose Sodium, Carnauba Wax, Magnesium Stearate, Food Red No.3 are contained as inactive ingredients..

(Efficacy)

Constipation,

Relief of the following symptoms accompanied with constipation:

Heaviness of head, a rush of blood, chap, skin eruption, loss of appetite, enlarged sensation of stomach, abnormal fermentation of intestine, hemorrhoids

(Dosage & Administration)

Each single dose is taken with water without chewing before going to bed.

Initial dose should be minimum and increase or decrease gradually depending on the efficacy observed.

<u>Age</u>	<u>Single dose</u>	<u>Number of dose/day</u>
Adults(over 15 years)	2 ~ 3 tablets	1 time
11 ~ 14 years	1 ~ 2 tablets	1 time
Below 11 years	Medication is not allowed.	

〈Caution for dosage and administration〉

- (1) Above dosage schedule should be followed strictly.
- (2) Medication for small children should be supervised by their parents.
- (3) As BEAULUCK is enteric-coated tablets, take without chewing or breaking tablets.
- (4) Do not take BEAULUCK within 1 hour after taking antacid or drinking milk.

(Caution for storage and handling)

- (1) Store at cool place with low humidity and no direct sunlight after closing tightly.
- (2) Keep out of the reach of the children.
- (3) Re-bottling of the tablets in other container is not allowed to avoid misuse or to maintain quality of medicine.

Manufacturer and Distributor:

KOKANDO PHARMACEUTICAL CO., LTD.

8-27, Nagasuhondori 2-Chome Amagasaki shi, Hyogo ken

JAPAN